

TYPE 2 AGNS WITH DOUBLE-PEAKED [O III] LINES: NARROW LINE REGION KINEMATICS OR MERGING SUPERMASSIVE BLACK HOLE PAIRS?

XIN LIU¹, YUE SHEN¹, MICHAEL A. STRAUSS¹, AND JENNY E. GREENE^{1,2}

Accepted to ApJ, November 3, 2009

ABSTRACT

We present a sample of 167 type 2 AGNs with double-peaked [O III] $\lambda\lambda 4959, 5007$ narrow emission lines, selected from the Seventh Data Release of the Sloan Digital Sky Survey. The double-peaked profiles can be well modeled by two velocity components, blueshifted and redshifted from the systemic velocity. Half of these objects have a more prominent redshifted component. In cases where the $H\beta$ emission line is strong, it also shows two velocity components whose line-of-sight (LOS) velocity offsets are consistent with those of [O III]. The relative LOS velocity offset between the two components is typically a few hundred km s^{-1} , larger by a factor of ~ 1.5 than the line full width at half maximum of each component. The offset correlates with the host stellar velocity dispersion σ_* . The host galaxies of this sample show systematically larger σ_* , stellar masses, and concentrations, and older luminosity-weighted mean stellar ages than a regular type 2 AGN sample matched in redshift, [O III] $\lambda 5007$ equivalent width and luminosity; they show no significant difference in radio properties. These double-peaked features could be due to narrow-line region kinematics, or binary black holes. The statistical properties do not show strong preference for or against either scenario, and spatially resolved optical imaging, spectroscopy, radio or X-ray followup are needed to draw firm conclusions.

Subject headings: black hole physics – galaxies: active – cosmology: observations – quasars: general – surveys

1. INTRODUCTION

Binary³ supermassive black holes (SMBHs) are possible outcomes of the hierarchical mergers of galaxies (e.g., Begelman et al. 1980; Milosavljević & Merritt 2001; Yu 2002). In one of the leading hypotheses, major mergers between galaxies are responsible for triggering nuclear starbursts and quasar activity (e.g., Hernquist 1989). Despite the success of the merger scenario in explaining much of the observed phenomenology of AGN statistics (e.g., Kauffmann & Haehnelt 2000; Volonteri et al. 2003; Wyithe & Loeb 2003; Hopkins et al. 2008; Shen 2009) and the cores of massive elliptical galaxies (e.g., Faber et al. 1997; Kormendy & Bender 2009), direct observational evidence for binary SMBHs is surprisingly scarce.

The fraction of binary quasars at separations of tens to hundreds of kpc (halo) scales, is $\lesssim 0.1\%$ at $1 \lesssim z \lesssim 5$ (e.g., Hennawi et al. 2006; Myers et al. 2008; Hennawi et al. 2009). On kpc (galactic) scales there are only a handful of unambiguous low-redshift cases known, in which both active SMBHs are detected in X-rays (NGC 6240 and Mrk 463, Komossa et al. 2003; Bianchi et al. 2008), or in the radio (3C 75, Owen et al. 1985). On sub-kpc scales, only one case is known (0402+379, Rodriguez et al. 2006) of a pair of BHs detected by VLBI with a projected separation of ~ 7 pc. There are several possible reasons that binary SMBHs are so rare compared to the expectations from the merger scenario: they spend most of their time at separations far below kpc scales and hence are difficult to resolve spatially; one or both of the SMBHs are heavily obscured; both BHs are rarely active at the same time, and dynamical differences between a single BH and a binary BH are not readily discernable at cosmolog-

ical distances.

Recently, Comerford et al. (2009a) conducted a systematic survey of type 2 AGNs in the DEEP2 galaxy sample at $\bar{z} \sim 0.6$ (Davis et al. 2003). They argue that 37 of the 107 AGNs in their sample are inspiralling binary SMBHs, based on significant velocity offsets between the narrow-line region (NLR) emission lines [O III] $\lambda\lambda 4959, 5007$ and the systemic redshift measured from stellar absorption features. In particular, two of these objects (one reported by Gerke et al. 2007) show double-peaked [O III] lines with velocity offsets of a few hundred km s^{-1} and spatial offsets of several kpc; they interpreted these as binary SMBHs when both BHs are active. Such double-peaked [O III] emission line features have also been seen in a few optically-selected AGNs (e.g., Heckman et al. 1981; Zakamska et al. 2003; Zhou et al. 2004). One advantage of using spectroscopy to find binary SMBHs is that it is not limited by spatial resolution, as long as the separation is not so small that the NLRs are no longer distinct. On the other hand, such spectral features may be due to biconical outflows or disk rotation around a single SMBH (e.g., Axon et al. 1998; Veilleux et al. 2001; Crenshaw et al. 2009). Here we focus on double-peaked *narrow line* objects in which the associated line emitting gas is on scales of ~ 100 pc to several kpc. Therefore in the binary SMBH scenario, our double-peaked sample would include objects whose two SMBHs along with their own NLR gas are still co-rotating in the galactic potential, well before forming a pc-scale binary under dynamical friction (e.g., Milosavljević & Merritt 2001). For double-peaked *broad line* objects which may involve outflows or disks on $\lesssim \text{pc}$ scales, see, e.g., Eracleous & Halpern (1994) and Strateva et al. (2003).

We have carried out a systematic search in the SDSS (York et al. 2000) spectroscopic database for emission line AGNs with double-peaked features or peculiar [O III] line shifts relative to the systemic redshift as measured from stellar absorption features. Here we report our initial results on a sample of 167 type 2 AGNs with double-peaked [O III] emis-

¹ Department of Astrophysical Sciences, Princeton University, Peyton Hall – Ivy Lane, Princeton, NJ 08544

³ Hubble Fellow, Princeton-Carnegie Fellow

³ We refer to both a pair of SMBHs in a Keplerian orbit and dual SMBHs at large separations where the galactic potential dominates as *binaries* unless otherwise noted.

TABLE 1
THE SAMPLE

SDSS Designation	Plate	Fiber	MJD	z	σ_*	FWHM _{[O III],1}	FWHM _{[O III],2}	$V_{[O III],1}$	$V_{[O III],2}$	$V_{H\beta,1}$	$V_{H\beta,2}$
J000249.07+004504.8	388	345	51793	0.0868	243	271	219	-361	168	-282	197
J000911.58-003654.7	388	148	51793	0.0733	203	231	220	-198	134	-196	129
J010750.48-005352.9	670	55	52520	0.5202	0	281	645	-546	37	-540	-113
J011659.59-102539.1	660	213	52177	0.1503	181	304	203	-159	145	-81	189

NOTE. — The full table is available in the electronic version of the paper. The subscripts “1” and “2” denote blueshifted and redshifted components. $V_{[O III]}$ and $V_{H\beta}$ denote velocity offsets relative to the systemic redshift measured over [O III] $\lambda\lambda 4959, 5007$ and H β . Velocities are in units of km s^{-1} . The entry reads zero if the quantity is unmeasurable. The typical uncertainty in σ_* is $\sim 15 \text{ km s}^{-1}$. For [O III], the typical statistical errors in the best-fit velocity offsets and FWHMs are $\sim 5 \text{ km s}^{-1}$ and $\sim 10 \text{ km s}^{-1}$. For H β , the statistical errors are generally a few times larger because H β is weaker than [O III].




FIG. 1.— Redshift distribution of the parent AGN and double-peaked samples.

sion lines⁴. We describe our sample selection in §2; the statistical properties of the sample are presented in §3, and we discuss our results and conclude in §4. A cosmology with $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, and $h = 0.7$ is assumed throughout.

2. DATA

2.1. The Sample

Our parent sample is the MPA-JHU SDSS DR7 galaxy sample⁵ drawn from the SDSS DR7 spectroscopic database (Abazajian et al. 2009); those include objects spectrally classified as galaxies by the *specBS* pipeline (Adelman-McCarthy et al. 2008) or quasars that are targeted as galaxies (Strauss et al. 2002; Eisenstein et al. 2001) and have redshifts $z < 0.7$. We adopt redshifts z and stellar velocity dispersions σ_* from the *specBS* pipeline. When unavailable from the pipeline, we measure σ_* using the direct-fitting algorithm described in Greene & Ho (2006) and Ho et al. (2009); we have checked that this code gives results consistent with those of *specBS*. We have checked the accuracy of the redshift by refitting the stellar continuum

⁴ After we submitted the paper, two similar statistical studies by Smith et al. (2009) and Wang et al. (2009) appeared on arXiv. Smith et al. (2009) focus on broad line AGNs with double-peaked [O III] lines selected from SDSS quasar spectra and find a similar fraction ($\sim 1\%$) of double-peaked objects that we found. Because of pipeline misclassification, their double-peaked sample also contains narrow line objects, some of which overlap with our catalog. Wang et al. (2009) studied a sample of 87 double-peaked narrow line AGNs of which 42 objects overlap with our sample.

⁵ <http://www.mpa-garching.mpg.de/SDSS/>

with galaxy templates (Liu et al. 2009), as well as comparing multiple epoch spectra of the same objects. The quoted redshift faithfully traces the stellar absorption features (which we take as the systemic redshift), with typical statistical errors of a few km s^{-1} and systematic errors of $\sim 10 \text{ km s}^{-1}$. Additional spectral and photometric properties such as emission-line fluxes (determined from equivalent widths and continuum fluxes), spectral indices, stellar masses, and concentration indices are taken from the MPA-JHU SDSS DR7 data product.

We select our AGN sample from this parent sample by the following criteria: 1) the rest-frame wavelength ranges $[4700, 5100] \text{ \AA}$ and $[4982, 5035] \text{ \AA}$ centered on the [O III] $\lambda 5007$ line have median signal-to-noise ratio (S/N) $> 5 \text{ pixel}^{-1}$ and bad pixel fraction $< 30\%$; 2) the [O III] $\lambda 5007$ line is detected at $> 5\sigma$ and has a rest-frame equivalent width (EW) $> 4 \text{ \AA}$; 3) the line flux ratio [O III] $\lambda 5007/\text{H}\beta > 3$ if $z > 0.33$, or the diagnostic line ratios [O III] $\lambda 5007/\text{H}\beta$ and [N II] $\lambda 6584/\text{H}\alpha$ lie above the theoretical upper limits for star-formation excitation from Kewley et al. (2001) on the BPT diagram (Baldwin et al. 1981) if $z < 0.33$. This procedure yields $\sim 14,300$ type 2 AGNs. We supplement this sample with ~ 400 type 2 quasars from Reyes et al. (2008) which are not included in the MPA-JHU data products. These additional type 2 quasars extend to somewhat higher redshifts ($z < 0.83$). Our final AGN sample includes 14,756 objects with high S/N and high spectral quality around the [O III] lines, suitable for the analysis that follows. The redshift distribution of this parent AGN sample is shown in Figure 1.

In many AGNs and starbursts, the narrow forbidden emission lines [O III] $\lambda\lambda 4959, 5007$ are known to have peculiarities such as extended blue wings, velocity offsets from systemic redshifts, and complex line profiles (e.g., Heckman et al. 1981; Whittle 1985a; Zakamska et al. 2003; Zhou et al. 2006; Komossa et al. 2008). We here focus on a more dramatic subset – those with double-peaked [O III] emission lines. A complete analysis of the nature and statistical properties of other peculiar [O III] line characteristics will be presented elsewhere.

The identification of such double-peaked objects requires that both [O III] $\lambda 4959$ and [O III] $\lambda 5007$ are better fit with two components rather than a single component. As the initial screening process we visually inspected the spectra of all the 14,756 AGNs and identified 167 objects with unambiguous double-peaked [O III] lines. We only include objects which have well-detected double peaks in both [O III] $\lambda 4959$ and [O III] $\lambda 5007$ with similar profiles; we do not include those with complex line profiles such as lumpy, winged, or multi-component features. We plot the redshift distribution of the double-peaked sample in Figure 1, and the SDSS images and




FIG. 2.— Example double-peaked AGNs. Shown here are SDSS *gri* color-composite images and host-subtracted spectra (data points in red and smoothed curve in black) along with our best fits (model in green and individual components in magenta) for the $H\beta$ -[O III] region. The vertical lines are drawn at the systemic redshift of each galaxy as measured from stellar absorption features.

spectra of three such examples in Figure 2. We list all the 167 double-peaked AGNs and their line measurements in Table 1. We caution that there could be some double-peaked narrow-line objects in SDSS DR7 missed by our selection if the pipeline got the redshifts wrong because of the peculiar line profiles, although this fraction is likely to be no larger than a few percent.

We measured the line properties of these double-peaked objects as follows. The galaxy continuum was subtracted using the best-fit template constructed from a linear combination of instantaneous starburst models of Bruzual & Charlot (2003) with ten different ages, as described in Liu et al. (2009). In the 11 objects for which the stellar continuum is too weak for measuring σ_* and for template fitting, a simple power-law model was used. The continuum-subtracted [O III] region ($\lambda\lambda 4930-5040\text{\AA}$) was fit by a pair of Lorentzian functions convolved with the measured instrumental resolution of the spectra ($\sigma \sim 65 \text{ km s}^{-1}$). The redshift and line width for each velocity component of [O III] $\lambda 4959$ and [O III] $\lambda 5007$

were forced to be the same. The flux ratio of [O III] $\lambda 5007$ to [O III] $\lambda 4959$ was allowed to vary (but we found it was always close to 3). In cases where $H\beta$ is measurable, we further fit a double-Lorentzian to the $H\beta$ region ($\lambda\lambda 4850-4880 \text{\AA}$), where the positions of the centroids were allowed to vary, but the widths of the two $H\beta$ components were fixed to the best-fit values for the two [O III] components. We then repeated this fitting process with a double-Gaussian model, and took as the best fit the model which had the smaller reduced χ^2 for each object. We visually inspected all the fits and verified that the line profiles are generally well reproduced by the model. For [O III], the typical statistical errors in the best-fit velocity difference between the two components v_{off} , line fluxes and FWHMs are $\sim 5 \text{ km s}^{-1}$, $\sim 5\%$, and $\sim 10 \text{ km s}^{-1}$, but we caution that the actual uncertainties are likely to be larger. For $H\beta$, the statistical errors are generally a few times larger because $H\beta$ is weaker than [O III].

2.2. Selection Completeness




FIG. 3.— Selection completeness as a function of median S/N , v_{off} , FWHM, total $[\text{O III}] \lambda 5007$ EW and the flux ratio between the two components, as determined from our Monte Carlo simulations. In the middle panel, black and blue lines are for the two components respectively.




FIG. 4.— Narrow-line region physical properties of the double-peaked components. Objects with stronger redshifted components are plotted as red open circles whereas those with stronger blueshifted components are plotted as blue plus signs. The Spearman correlation coefficient and the probability of null correlation are labelled in each plot for the whole (black), the stronger-blueshifted (blue), and the stronger-redshifted (red) samples, respectively. (a). Ionization parameter (indicated by the flux ratio between $[\text{O III}] \lambda 5007$ and $\text{H}\beta$) of the blueshifted component versus that of the redshifted component. (b). Line width ratio of the double-peaked components versus Balmer decrement (as a measure of reddening). (c). Electron density (inferred from the flux ratio between the $[\text{S II}] \lambda \lambda 6717, 6731$ doublet) of the blueshifted component versus that of the redshifted component.

Our selection of double-peaked objects was done by eye and is by no means objective. To gain some sense of our completeness we performed Monte Carlo simulations. We generated mock spectra of double-peaked objects in the $\text{H}\beta$ - $[\text{O III}]$ region with random distributions of continuum S/N , EW, velocity offset, and FWHM of the two components of $[\text{O III}]$. We randomly assigned a Gaussian or Lorentzian profile to each line and broadened the spectrum using an instrumental Gaussian broadening of $\sigma = 65 \text{ km s}^{-1}$. In this way we generated 2,000 mock spectra of double-peaked objects and mixed them with 18,000 mock spectra of single-peaked objects with random spectral properties. We then visually inspected the 20,000 mock spectra and identified double-peaked objects in the same way as we did for the real sample. We show in Figure 3 the selection completeness as a function of median S/N , v_{off} , FWHM, total $[\text{O III}] \lambda 5007$ EW, and flux (EW) ratio between the two components f_{EW} . In general the completeness increases with increasing v_{off} , $[\text{O III}] \lambda 5007$ EW, and f_{EW} , and decreases with increasing FWHM, as expected from the way our double-peaked objects are identified. As a result there is a selection bias against objects with small v_{off} and large FWHM that we discuss later. In addition we are very incomplete for objects with $v_{\text{off}} \lesssim 200 \text{ km s}^{-1}$ partly due to the spectral resolution limit⁶. On the other hand, we find that the false de-

tection rate is very low ($< 2\%$ of the double-peaked sample) – probably an advantage of visual inspection over automated algorithms. We note that the completeness estimated this way is not the *absolute* completeness, which depends on the actual underlying distributions of all relevant properties. Nevertheless, these completeness estimates are useful to correct for selection biases. For example, when other properties are fixed, double-peaked objects with larger EWs are easier to identify, by a relative amount that can be determined from our simulation results.

After accounting for the dependence of selection completeness on $[\text{O III}]$ EW, we find tentative evidence that the fraction of double-peaked objects in the underlying AGN sample depends on $[\text{O III}] \lambda 5007$ luminosity ($L_{[\text{O III}]}$): it increases by a factor of ~ 2 from $L_{[\text{O III}]} \sim 10^{7.0} L_{\odot}$ to $L_{[\text{O III}]} \sim 10^{8.5} L_{\odot}$, and has no clear dependence on redshift in the range probed. In most cases the two velocity components are redshifted and blueshifted with respect to the systemic redshift (except that in a few cases, one component is apparently coincident with the systemic velocity within the errors), and about half of them have more prominent redshifted components – a subset which may be more unusual since objects with outflows may tend to have more prominent blueshifted components (see below). In cases where $\text{H}\beta$ is measurable, it also shows a double-peaked feature with a similar velocity offset as that of $[\text{O III}]$. Hereafter we use v_{off} measured from $[\text{O III}]$ in our analysis as it is more robust than that from $\text{H}\beta$. As illustrated in Figure 4(a), both velocity components have excitation diagnostic line ra-

⁶ For instance, the object SDSS J100043.13+020637.2, with $v_{\text{off}} \sim 150 \text{ km s}^{-1}$ reported in Comerford et al. (2009b) is not included in our sample based on the SDSS spectrum alone.




FIG. 5.— Host galaxy properties (stellar velocity dispersion σ_* , stellar mass M_* , spectral indices $D_n(4000)$ and $H\delta_A$ indicating the luminosity-weighted mean stellar age and the post-starburst fraction, and r -band concentration index $R_{90,r}/R_{50,r}$) of the double-peaked sample (red) compared to the control sample (black). The control sample is randomly drawn from the parent AGN sample to have identical redshift and $[O\text{ III}]\lambda 5007$ EW distributions as the double-peaked sample. For each distribution the median is marked with a vertical line.

tios $H\beta/[O\text{ III}]$ characteristic of AGNs⁷. The FWHM of each velocity component is typically smaller than v_{off} by a factor of ~ 1.5 .

3. STATISTICAL PROPERTIES

3.1. Host Galaxy Properties

Are there any characteristics of the host galaxies of the double-peaked objects that are different from ordinary AGNs? The double-peaked sample is not selected uniformly as we demonstrated in §2.2. To account for selection biases, we construct a control sample from the underlying AGN sample with identical redshift, $[O\text{ III}]\lambda 5007$ EW and luminosity distributions as the double-peaked sample. We compare their host-galaxy properties, including stellar mass and velocity dispersion, SDSS colors, stellar population parameters $D_n(4000)$ and $H\delta_A$ (indicating the luminosity-weighted mean stellar age and the post-starburst fraction in the past ~ 0.1 – 1 Gyr, respectively; e.g., Kauffmann et al. 2003), intrinsic extinction (estimated by the Balmer decrement), effective radius, concentration (characterized by, $R_{90,r}/R_{50,r}$, the ratio of radii that contain 90% and 50% of the r -band Petrosian flux), and inclination (estimated by the ratio between the major and minor isophotal axis). As shown in Figure 5, the double-peaked sample has systematically larger stellar velocity dispersions and masses, older mean stellar ages (and redder colors), smaller fractions of post-starburst populations in the past 0.1–1 Gyr, and higher concentrations than the control sample. The difference in the centroids of the two distributions are roughly half of the width of the distributions. Kolmogorov-Smirnov (KS) tests show that the probabilities that the two samples are drawn from the same distribution are $P_{\text{KS}} < 10^{-3}$. All other properties of the host galaxies we examined were indistinguishable with KS probabilities $P_{\text{KS}} > 10^{-1}$. We also match the double-peaked and the control samples with the FIRST (Becker et al. 1995; White et al. 1997) and ROSAT (Voges et al. 1999, 2000) surveys and find that they contain very similar fractions of matches with both surveys ($\sim 35\%$ with FIRST and $\sim 2\%$ with ROSAT). The subset that has a more prominent redshifted component is indistinguishable from the rest of the double-peaked sample in terms of the properties studied above.

3.2. Correlations between Line Properties

There are several apparent correlations among the dynamical properties and the NLR physical conditions of the

double-peaked sample. Figures 4(a) and 4(c) illustrate that the two velocity components have similar ionization parameters (as indicated by the diagnostic line ratio $H\beta/[O\text{ III}]$) and electron densities (measured from the ratio of the doublet $[S\text{ II}]\lambda\lambda 6717, 6731$ for 107 objects in the sample which have low enough redshifts to have $[S\text{ II}]$ coverage in the SDSS spectra and whose inferred electron densities of both components fall in the range of $[10, 10^4]\text{ cm}^{-3}$). Figures 6(a)–(d) show that v_{off} is correlated with both σ_* and the line FWHMs of the two components, and the line FWHMs are correlated with each other. In all these four relations, the correlation for objects with more prominent redshifted components appears somewhat stronger than that for the rest. There is some selection bias at the low v_{off} versus large FWHM corner, where the double-peaked feature is difficult to identify. However, our Monte-Carlo simulations show no selection bias against objects with high v_{off} and small FWHMs (§2.2). In particular, the correlation between v_{off} and σ_* suggests that the double-peaked line-emitting regions are on galactic scales where the galactic potential dominates the bulk kinematics. Furthermore, Figures 6(e) and 6(f) show that both the $[O\text{ III}]$ line flux ratio and the line width ratio of the two velocity components are anti-correlated with the ratio of their LOS velocity offsets relative to the systemic redshift (see also Wang et al. 2009), which most likely results from momentum conservation.

4. DISCUSSION

4.1. NLR Kinematics

The double-peaked feature may result from particular NLR geometries such as biconical outflows⁸ or rotating disks on kpc scales. In these scenarios, there is only one SMBH and the observed blueshifted and redshifted emission-line peaks arise from NLR gas moving towards and away from us. Certain nearby Seyfert galaxies known to have biconical outflows show double-peaked emission when spectra are taken over the whole galaxy. Such prototypical cases include NGC 1068 (e.g., Axon et al. 1998) and NGC 3079 (e.g., Duric & Seaquist 1988). It is very likely that there are some such sources in our sample. Several objects in our sample show clear galactic disks in their SDSS images. Here the correlation between line width and velocity shift naturally arises, as the bulge gravitational potential well influences the motion of NLR gas. The correlation between the diagnostic line ratio $[O\text{ III}]/H\beta$ of the two components shown in Figure 4(a) may

⁷ For objects with $z < 0.3$, the $[N\text{ II}]$ and $H\alpha$ lines are visible; they appear double-peaked as well, although the different components often are blended with one another.

⁸ In principle, a NLR with inflowing gas and dust can also produce the observed features, but this scenario seems less favored considering the difficulty of removing high angular momentum given the large v_{off} (e.g., Heckman et al. 1981).




FIG. 6.— Dynamical properties of the double-peaked components. Objects with stronger redshifted components are plotted as red open circles whereas those with stronger blueshifted components are plotted as blue plus signs. The Spearman correlation coefficient and the probability of null correlation are labelled in each plot for the whole (black), the stronger-blueshifted (blue), and the stronger-redshifted (red) samples, respectively. (a)–(c). stellar velocity dispersion σ_* , FWHM of the blueshifted component, and FWHM of the redshifted component versus velocity offset between the double-peaked components. (d). FWHM of the blueshifted component versus that of the redshifted component. (e) Velocity-offset ratio of the double-peaked components versus the flux ratio. (f). Velocity-offset ratio of the double-peaked components versus the FWHM ratio.

be explained if the NLR gas of both velocity components is ionized by a single SMBH.

Can all the objects in our double-peaked sample be ascribed to NLR geometry? Outflows and inflows of NLR gas have long been invoked to explain several commonly observed features in NLR high ionization lines (such as [O III]), including extended wings (usually blueshifted from the systemic redshift), line asymmetries, and broader line widths than those of low ionization lines (e.g., Heckman et al. 1981; Penston et al. 1984; Rodríguez-Ardila et al. 2006). In the cases of NGC 1068 (e.g., Axon et al. 1998) and Mrk 78 (e.g., Heckman et al. 1981; Pedlar et al. 1989; Whittle & Wilson 2004) known to have biconical outflows, the velocity splitting relative to the systemic redshift is larger on the blueshifted component, which is also brighter and has a larger width than the redshifted component. The statistical significance of these trends is unclear, however (e.g., Whittle et al. 1988). Unlike these, our double-peaked sample has comparable line widths in the blueshifted and redshifted components as shown in Figure 6(d); half of our double-peaked objects have a more prominent redshifted component than the blueshifted component. In addition, our sample has consistent v_{off} in H β and in [O III] whenever H β is measurable, with no significant ionization stratification as expected in NLR outflows. These possible discrepancies may arise because our sample has appreciably larger [O III] luminosity than these previous low redshift comparison samples, and these properties may be a function of lu-

minosity or redshift or both; alternatively, they may suggest that not all objects in our double-peaked sample exhibit NLR outflows.

Further possible clues come from host galaxy and line properties. The FIRST detected fraction for the double-peaked objects is comparable to that of the control sample. The host-galaxy inclination distribution shows no significant difference either, whereas our selection of a large LOS v_{off} for the double components would bias towards either edge-on or face-on inclinations for the rotating disk or bipolar outflow populations, unless the outflow orientation is not closely linked to the angular momentum axis of the host galaxies. Radio outflows in nearby Seyfert galaxies and low-ionization AGNs indeed appear to orient randomly with respect to the host galaxy axes (e.g., Ulvestad & Wilson 1984; Gallimore et al. 2006). In addition, while a correlation between v_{off} and line FWHM is observed in NLR outflows (e.g., Komossa et al. 2008), the ratio of velocity shift to line width is of order unity, unlike the ratio of ~ 1.5 seen in our double-peaked sample. Finally, as shown in Figure 4(b), there is no obvious correlation in our sample between the Balmer decrement and the line-width ratio between the two components of the double-peaked feature (as a proxy for asymmetry). Such a correlation might exist if the blueshifted and redshifted NLR gas components are intrinsically symmetric and the apparent asymmetry results from extinction. Heckman et al. (1981) observed a correlation between the asymmetry of a single line profile (which, how-

ever we caution, is different from our asymmetry indicator – the line width ratio of the double-peaked components) and Balmer decrement in a sample of 36 nearby Seyferts thought to contain outflows, although later studies on nearby Seyferts (e.g., Whittle 1985b) found no strong correlations.

4.2. Merging SMBH Pairs

Another explanation for the double-peaked features is merging SMBH pairs. In this scenario, both BHs, along with their own NLR gas (with scales of order hundreds of pc), are co-rotating in the galactic potential on $\gtrsim$ kpc scales, well before dynamical friction causes their orbit to decay and form a gravitationally bounded compact binary (e.g., Milosavljević & Merritt 2001). The $v_{\text{off}}-\sigma_*$ and $v_{\text{off}}\text{-FWHM}$ correlations in this scenario would naturally arise from dynamics. The comparable numbers of objects with more prominent redshifted and blueshifted components and the null correlation between the Balmer decrement and the double-peak asymmetry may both be accommodated if the two emission components are associated with two SMBHs. The similar distribution of host-galaxy inclination angles to the control sample could be explained, given that there is no preference for the rotation axis of galaxies in the merging SMBH-pair scenario.

On the other hand, the correlation between the diagnostic line ratio $[\text{O III}]/\text{H}\beta$ of the two components shown in Figure 4(a) may be difficult to explain if the NLR gas of each component is primarily ionized by its own BH. In addition, the higher galaxy concentration for objects with double peaks (Figure 5) seems inconsistent with these objects being in the early stage of mergers, although we may be biased to more compact objects by the necessity to get both merging objects within the 3-arcsec fiber aperture. The older mean stellar ages and the smaller post-starburst fraction in the past 0.1–1 Gyr also seem counterintuitive, if there is excess star formation in mergers. However, these differences might also result from the mass-age correlation (e.g., Kauffmann et al. 2003) given that the median σ_* of the double-peaked sample is larger than that of the control sample (Figure 5), or if the associated star formation is on a different time scale from what is probed by the $D_n(4000)$ and $\text{H}\delta_A$ indices adopted here, or it is suppressed by AGN activity.

In view of the above arguments, we conclude that our double-peaked sample may contain a population of NLR outflows or rotating disks, as well as objects that are merging SMBHs; The statistical properties of the sample do not show any strong preference for or against either scenario. In Figure 2 we show several intriguing cases. The first example (SDSSJ114642.5+511029.9) is a merging system of which the off-centered galaxy is partially covered by the 3-arcsec diameter fiber. We have checked that the two galaxies are both detected in the K_s band by 2MASS (Skrutskie et al. 2006), and have found evidence that both are AGNs using spatially resolved longslit spectroscopy, the results of which will be presented elsewhere. Only 13 of the 167 objects in our double-peaked sample show possible evidence for two cores in the SDSS images within the 3-arcsec diameter of the spectro-

scopic fibers. The second object (SDSSJ122709.8+124854.5) does not have a resolved double core in its SDSS image, and its emission lines have well separated peaks, symmetrically shifted from the systemic velocity, and the redshifted component is more prominent. High resolution HST imaging may help reveal potential double cores unresolved in SDSS images (e.g., Comerford et al. 2009b). For example, one of our double-peaked objects (SDSSJ130128.8–005804.3, at $z = 0.2455$) shows no double cores in its SDSS image, but Zakamska et al. (2006) used HST imaging to show that it consists of two galaxies with a projected separation of 1.3 arcsec (corresponding to 5.0 kpc) which well fits into the SDSS 3-arcsec diameter fiber. We will need spatially resolved spectroscopy to determine whether both of the galaxies are AGN. The last object in Figure 2 has a strong component coincident with the systemic redshift and a weaker blueshifted component and an even weaker redshifted component (not fitted), which could arise from a classic NLR region plus bipolar outflows. Spatially-resolved optical imaging, spectroscopy, radio and/or X-ray followup are still needed to help draw firm conclusions on the nature of these double-peaked narrow line objects.

We thank J. Krolik and W. Voges for helpful comments, and an anonymous referee for a careful and useful report that improves the paper. X.L., Y.S., and M.A.S. acknowledge the support of NSF grant AST-0707266. Support for J.E.G. was provided by NASA through Hubble Fellowship grant HF-01196 awarded by the STSI, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org/>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

Facilities: Sloan

REFERENCES

- Abazajian, K. N., et al. 2009, *ApJS*, 182, 543
- Adelman-McCarthy, J. K., et al. 2008, *ApJS*, 175, 297
- Axon, D. J., Marconi, A., Capetti, A., Maccetto, F. D., Schreier, E., & Robinson, A. 1998, *ApJ*, 496, L75
- Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, *PASP*, 93, 5
- Becker, R. H., White, R. L., & Helfand, D. J. 1995, *ApJ*, 450, 559
- Begelman, M. C., Blandford, R. D., & Rees, M. J. 1980, *Nature*, 287, 307
- Bianchi, S., Chiaberge, M., Piconcelli, E., Guainazzi, M., & Matt, G. 2008, *MNRAS*, 386, 105
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000

- Comerford, J. M., et al. 2009a, *ApJ*, 698, 956
- Comerford, J. M., Griffith, R. L., Gerke, B. F., Cooper, M. C., Newman, J. A., Davis, M., & Stern, D. 2009b, *ApJ*, 702, L82
- Crenshaw, D. M., Schmitt, H. R., Kraemer, S. B., Mushotzky, R. F., & Dunn, J. P. 2009, *ApJ* submitted
- Davis, M., et al. 2003, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 4834, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, ed. P. Guhathakurta, 161–172
- Duric, N., & Seaquist, E. R. 1988, *ApJ*, 326, 574
- Eisenstein, D. J., et al. 2001, *AJ*, 122, 2267
- Eracleous, M., & Halpern, J. P. 1994, *ApJS*, 90, 1
- Faber, S. M., et al. 1997, *AJ*, 114, 1771
- Gallimore, J. F., Axon, D. J., O’Dea, C. P., Baum, S. A., & Pedlar, A. 2006, *AJ*, 132, 546
- Gerke, B. F., et al. 2007, *ApJ*, 660, L23
- Greene, J. E., & Ho, L. C. 2006, *ApJ*, 641, 117
- Heckman, T. M., Miley, G. K., van Breugel, W. J. M., & Butcher, H. R. 1981, *ApJ*, 247, 403
- Hennawi, J. F., et al. 2009, *ArXiv e-prints*
- . 2006, *AJ*, 131, 1
- Hernquist, L. 1989, *Nature*, 340, 687
- Ho, L. C., Greene, J. E., Filippenko, A. V., & Sargent, W. L. W. 2009, *ApJS*, 183, 1
- Hopkins, P. F., Hernquist, L., Cox, T. J., & Kereš, D. 2008, *ApJS*, 175, 356
- Kauffmann, G., & Haehnelt, M. 2000, *MNRAS*, 311, 576
- Kauffmann, G., et al. 2003, *MNRAS*, 346, 1055
- Kewley, L. J., Dopita, M. A., Sutherland, R. S., Heisler, C. A., & Trevena, J. 2001, *ApJ*, 556, 121
- Komossa, S., Burwitz, V., Hasinger, G., Predehl, P., Kaastra, J. S., & Ikebe, Y. 2003, *ApJ*, 582, L15
- Komossa, S., Xu, D., Zhou, H., Storchi-Bergmann, T., & Binette, L. 2008, *ApJ*, 680, 926
- Kormendy, J., & Bender, R. 2009, *ApJ*, 691, L142
- Liu, X., Zakamska, N. L., Greene, J. E., Strauss, M. A., Krolik, J. H., & Heckman, T. M. 2009, *ApJ*, 702, 1098
- Milosavljević, M., & Merritt, D. 2001, *ApJ*, 563, 34
- Myers, A. D., Richards, G. T., Brunner, R. J., Schneider, D. P., Strand, N. E., Hall, P. B., Blomquist, J. A., & York, D. G. 2008, *ApJ*, 678, 635
- Owen, F. N., O’Dea, C. P., Inoue, M., & Eilek, J. A. 1985, *ApJ*, 294, L85
- Pedlar, A., Meaburn, J., Axon, D. J., Unger, S. W., Whittle, D. M., Meurs, E. J. A., Guerrine, N., & Ward, M. J. 1989, *MNRAS*, 238, 863
- Penston, M. V., Fosbury, R. A. E., Boksenberg, A., Ward, M. J., & Wilson, A. S. 1984, *MNRAS*, 208, 347
- Reyes, R., et al. 2008, *AJ*, 136, 2373
- Rodriguez, C., Taylor, G. B., Zavala, R. T., Peck, A. B., Pollack, L. K., & Romani, R. W. 2006, *ApJ*, 646, 49
- Rodríguez-Ardila, A., Prieto, M. A., Viegas, S., & Gruenwald, R. 2006, *ApJ*, 653, 1098
- Shen, Y. 2009, *ApJ*, 704, 89
- Skrutskie, M. F., et al. 2006, *AJ*, 131, 1163
- Smith, K. L., Shields, G. A., Bonning, E. W., McMullen, C. C., & Salvander, S. 2009, *ArXiv e-prints* 0908.1998
- Strateva, I. V., et al. 2003, *AJ*, 126, 1720
- Strauss, M. A., et al. 2002, *AJ*, 124, 1810
- Ulvestad, J. S., & Wilson, A. S. 1984, *ApJ*, 285, 439
- Veilleux, S., Shopbell, P. L., & Miller, S. T. 2001, *AJ*, 121, 198
- Voges, W., et al. 1999, *A&A*, 349, 389
- . 2000, *IAU Circ.*, 7432, 3
- Volonteri, M., Haardt, F., & Madau, P. 2003, *ApJ*, 582, 559
- Wang, J., Chen, Y., Hu, C., Mao, W., Zhang, S., & Bian, W. 2009, *ArXiv e-prints*
- White, R. L., Becker, R. H., Helfand, D. J., & Gregg, M. D. 1997, *ApJ*, 475, 479
- Whittle, M. 1985a, *MNRAS*, 213, 1
- . 1985b, *MNRAS*, 213, 33
- Whittle, M., Pedlar, A., Meurs, E. J. A., Unger, S. W., Axon, D. J., & Ward, M. J. 1988, *ApJ*, 326, 125
- Whittle, M., & Wilson, A. S. 2004, *AJ*, 127, 606
- Wyithe, J. S. B., & Loeb, A. 2003, *ApJ*, 595, 614
- York, D. G., et al. 2000, *AJ*, 120, 1579
- Yu, Q. 2002, *MNRAS*, 331, 935
- Zakamska, N. L., et al. 2003, *AJ*, 126, 2125
- . 2006, *AJ*, 132, 1496
- Zhou, H., Wang, T., Yuan, W., Lu, H., Dong, X., Wang, J., & Lu, Y. 2006, *ApJS*, 166, 128
- Zhou, H., Wang, T., Zhang, X., Dong, X., & Li, C. 2004, *ApJ*, 604, L33